



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

determine whether infection was as abundant under such a cover as in the open. Infection was fully as abundant. Conclusions could not be satisfactorily drawn, however, because it was found that the mesh of screen used was large enough under the conditions to allow the entrance of a large number of very small insects. The conditions of the experiment were, therefore, revised this year with the hope of making them crucial. Twelve-mesh wire screen cylinders, 15 cm. in diameter and 30 cm. long, were constructed to enclose parts of single branches. To prevent contact with the branches to be enclosed, cords were run four times transversely through each cylinder. Some of the cylinders were slipped into closely fitting sleeves of fine bolting cloth (124 threads per linear inch). For durability, the bolting cloth was sewed into canvas which formed the ends of the sleeves, covering the rough ends of the wire, and extending past the wire about 14 cm., sufficiently to permit secure tying. The ends of other cylinders were similarly covered with canvas and the exposed part of the wire was painted with a mixture of tangle-foot and benzene. The cylinders were slipped over branches including either last year's terminal shoots or bearing wood. In the former case the cylinders were extended far enough past the ends of the branches to allow for this year's terminal growth. All this was done before any of the blossoms opened, at the beginning of the pink-bud stage. The cylinders treated with tangle-foot were repainted frequently enough to maintain a sticky surface. No insects and no traces of them of any sort were found in any of the cages with one exception.⁸

There were ten cylinders enclosing flowering wood. Flowers in two of these cages blighted, as was shown by their appearance and as was verified by microscopic examination. The blight evidently entered through the calyx. Blossom blight was not abundant this season in this orchard.

Forty cylinders, twenty of the bolting cloth

⁸ One cylinder was accidentally allowed to dry. Two insects were found in it. The shoot was not blighted.

and twenty of the tangle-foot type, were used to include terminal growth. Of this year's shoots four in the bolting cloth cylinders and eight in the tangle-foot cylinders blighted, a total of twelve in forty, or thirty per cent. This was practically the same proportion of blighted terminal shoots as prevailed among the unenclosed shoots, as was shown by a count of a thousand terminal shoots on these and adjacent trees of the same variety and age.

From the above facts it appears that there must have been some agency of dispersal other than insects, and that insects were not even of primary importance as carriers. The only tenable hypothesis is that wind was the chief agent of transmission. Supporting evidence for this conclusion is found in two facts: (1) in the lack of insects in the orchard in sufficient numbers to account for the large amount of twig blight and (2) in the entire absence of insects from exuding cankers, whence they might receive their initial contamination. In three years' close observation at blooming time one of the authors (Ruth) has not observed a single case of insect visitation to exuding cankers. Aphids were entirely absent through the season of floral and twig infection. Leafhoppers became evident in rather small numbers only after the period of infection had passed. No other insects were present in sufficient numbers to be considered primary agents.

F. L. STEVENS,
W. A. RUTH,
C. S. SPOONER

UNIVERSITY OF ILLINOIS

SCIENCE

A Weekly Journal devoted to the Advancement of
Science, publishing the official notices and pro-
ceedings of the American Association for
the Advancement of Science

Published every Friday by

THE SCIENCE PRESS

LANCASTER, PA.

GARRISON, N. Y.

NEW YORK, N. Y.

Entered in the post-office at Lancaster, Pa., as second class matter